

Heaviside 1850-1925 Farad, Maxwell, Ampere

Math 231, Thu 22-Apr-2021 -- Thu 22-Apr-2021
Differential Equations and Linear Algebra
Spring 2021

Thursday, April 22nd 2021

Due:: HC05 due at 11 pm

Other calendar items

Thursday, April 22nd 2021

Wk 12, Th

Topic:: Heaviside unit step fn

Heaviside unit step

- its Laplace transform

original

right-shifted version

- use in writing other piecewise-defined functions

square pulse

sine function switched on at time $\pi/2$

Q: Which do we mean? Is the "shift" only in H , or in both functions?

$H(t - \pi/2) * \sin(t - \pi/2)$

$H(t - \pi/2) * \sin(t)$

ramp function


saw-tooth function

- Shift theorems

$$H(t) = \begin{cases} 0, & \text{if } t < 0 \\ 1, & \text{if } t > 0 \end{cases} \text{ piecewise-defined function}$$

Ex.) $\mathcal{L}\{H(t)\} = \int_0^\infty e^{-st} H(t) dt = \int_0^\infty e^{-st} \cdot 1 dt$

$$= \mathcal{L}\{1\} = \frac{1}{s}.$$



$$\mathcal{L}\{m H(t-c)\} = \int_0^\infty e^{-st} \cdot \underbrace{m H(t-c)}_{=0 \text{ on } 0 < t < c} dt$$

$$= m \text{ on } t > c$$

$$= \int_c^\infty e^{-st} \cdot m dt$$


$$= m \int_c^\infty e^{-st} dt = -\frac{m}{s} \left[e^{-st} \right]_c^\infty$$

$$\text{if } s > 0 = -\frac{m}{s} \left(0 - e^{-sc} \right) = m \cdot \frac{e^{-sc}}{s}.$$



Continuing our table

<u>time side</u>	<u>freq. side</u>
$H(t-c)$	e^{-sc}/s
$H(t-c) f(t-c)$	$e^{-sc} \cdot \mathcal{L}\{f(t)\}(s)$

Using $H(t)$ — name for particular PW-defined fn — in writing other PW-defined fns.




$$= \begin{cases} 0, & t < c \\ 1, & c < t < d \\ 0, & t > d \end{cases}$$



$$= H(t-c) - H(t-d)$$


Ex.) Ramp fn.




$$= \begin{cases} 0, & t < c \\ \frac{m}{d-c}(t-c), & c < t < d \\ m, & t > d \end{cases}$$

slope = $\frac{m}{d-c}$, passes thru $(c, 0)$

$$y - 0 = \frac{m}{d-c}(x - c)$$




Exercise?
Write this fn. using $H(t)$




Consider now that I might want $\mathcal{L}\{f(t)\}$
where $f(t)$ is a delayed sine fn.; say delayed by $\pi/2$.

Let's first get straight whether we mean

1. $H(t - \pi/2) \cdot \sin t$



2. $H(t - \pi/2) \sin(t - \pi/2)$



Shift theorem

$$\mathcal{L}\{H(t-c)f(t-c)\} = \int_0^\infty e^{-st} H(t-c)f(t-c) dt$$

$$= \int_c^\infty e^{-st} f(t-c) dt$$

Using substitution rule from Calculus

$$\text{Let } \tau = t - c \quad \left(\text{so } d\tau = dt \right)$$

$$= \int_0^\infty e^{-s(\tau+c)} f(\tau) d\tau$$

$$= e^{-sc} \left(\int_0^\infty e^{-s\tau} f(\tau) d\tau \right)$$

$$= \mathcal{L}\{f(t)\}$$

$$= e^{-sc} \cdot \mathcal{L}\{f(t)\}$$

Interpretation 2 above, if transformed

$$\begin{aligned} \mathcal{L}\{H(t - \pi/2) \sin(t - \pi/2)\} &= e^{-s \cdot \pi/2} \cdot \mathcal{L}\{\sin(t)\} \\ &= e^{-s \cdot \pi/2} \cdot \frac{1}{s^2 + 1} \end{aligned}$$

MATH 231

Laplace transform shift theorems

There are **two** results/theorems establishing connections between shifts and exponential factors of a function and its Laplace transform.

Theorem 1: If $f(t)$ is a function whose Laplace transform $\mathcal{L}[f(t)](s) = F(s)$, then

- A. $\mathcal{L}[e^{at}f(t)](s) = F(s-a)$, and
- B. $\mathcal{L}[H(t-a)f(t-a)](s) = e^{-as}F(s)$.

Neither of these theorems is strictly necessary for computing Laplace transforms—i.e., when going from the time domain function $f(t)$ to its frequency domain counterpart $\mathcal{L}[f(t)](s)$. Such transforms can be computed directly from the definition of Laplace transform $\mathcal{L}[f(t)](s) = \int_0^\infty e^{-st}f(t)dt$.

Example 1:

We compute

- (a) $\mathcal{L}[te^{2t}](s)$, and
- (b) $\mathcal{L}[H(t-3)e^{t-3}]$

directly from the definition.

For part (a),

$$\begin{aligned}\mathcal{L}[te^{2t}](s) &= \int_0^\infty e^{-st}te^{2t}dt = \int_0^\infty te^{-(s-2)t}dt = \int_0^\infty te^{-st}dt \Big|_{s \mapsto s-2} = \mathcal{L}[t](s-2) \\ &= \frac{1}{s^2} \Big|_{s \mapsto s-2} = \frac{1}{(s-2)^2}.\end{aligned}$$

For part (b),

$$\begin{aligned}\mathcal{L}[H(t-3)e^{t-3}] &= \int_0^\infty e^{-st}H(t-3)e^{t-3}dt = \int_3^\infty e^{-st}e^{t-3}dt \\ &= \int_0^\infty e^{-s(u+3)}e^u du \quad (\text{by substitution: } u = t-3) \\ &= e^{-3s} \int_0^\infty e^{-su}e^u du = e^{-3s} \int_0^\infty e^{-st}e^t dt \quad (\text{the name of the variable of integration is immaterial}) \\ &= e^{-3s} \mathcal{L}[e^t] = e^{-3s} \frac{1}{s-1}.\end{aligned}$$

■

Using shift theorems for inverse Laplace transforms

It is in finding *inverse* Laplace transforms where Theorems A and B are indispensable.

Example 2:

Find the inverse Laplace transform for each of the functions

$$(a) \frac{se^{-2s}}{s^2 + 9}$$

$$(b) \frac{3}{(s + 1)^3}$$

$$(c) \frac{2s}{s^2 - 4s + 5}$$

Our function in part (a) has an exponential factor, much like in Theorem B. Here,

$$e^{-2s} \frac{s}{s^2 + 9} = e^{-2s} F(s), \quad \text{where} \quad F(s) = \frac{s}{s^2 + 9} = \mathcal{L}[\cos(3t)](s).$$

Thus,

$$\mathcal{L}^{-1}\left[e^{-2s} \frac{s}{s^2 + 9}\right](t) = H(t - 2) \cos(3(t - 2)).$$

The function in part (b) does not look like an entry in the Laplace transform table I provide:

http://www.calvin.edu/~scofield/courses/m231/F14/table_of_Laplace_transforms.pdf

It is, in fact, a modified version of the table entry $n!/s^{n+1}$ with $n = 2$ but shifted left 1 unit, i.e.,

$$\frac{3}{(s + 1)^3} = \frac{3}{s^3} \Big|_{s \mapsto s+1} = \frac{3}{2} \cdot \frac{2!}{s^3} \Big|_{s \mapsto s-(-1)}.$$

Since

$$\mathcal{L}^{-1}\left[\frac{3}{2} \cdot \frac{2!}{s^3}\right](t) = \frac{3}{2} \mathcal{L}^{-1}\left[\frac{2!}{s^3}\right](t) = \frac{3}{2} t^2,$$

it follows from Theorem A that

$$\mathcal{L}^{-1}\left[\frac{3}{(s + 1)^3}\right](t) = \mathcal{L}^{-1}\left[\frac{3}{2} \cdot \frac{2!}{s^3} \Big|_{s \mapsto s-(-1)}\right](t) = \frac{3}{2} t^2 e^{-t}.$$

The function in part (c) also does not look like an entry in the table of Laplace transforms found at the link above. The denominator is, in fact, an **irreducible quadratic (over the reals)**, having no real roots. But a quadratic has a parabolic graph, and any parabola may be obtained from the graph of $y = x^2$ via a sequence of shifts and stretches. We can find the shift involved through completing the square:

$$s^2 - 4s + 5 = s^2 - 4s + 4 + 1 = (s - 2)^2 + 1,$$

which means the graph of $s^2 - 4s + 5$ is the same as the graph of $s^2 + 1$ but shifted 2 units to the right. To use Theorem A, we need *all* instances of s to be similarly shifted, so we write

$$\frac{2s}{s^2 - 4s + 5} = \frac{2s}{(s - 2)^2 + 1} = \frac{2(s - 2 + 2)}{(s - 2)^2 + 1} = \frac{2(s - 2) + 4}{(s - 2)^2 + 1} = \frac{2s + 4}{s^2 + 1} \Big|_{s \mapsto s-2}.$$

[Take a moment to plot, together, the functions $2x/(x^2 - 4x + 5)$ and $(2x + 4)/(x^2 + 1)$. Observe that the graph of the former is identical to that of the latter, except shifted right 2 units.] Since

$$\mathcal{L}^{-1}\left[\frac{2s + 4}{s^2 + 1}\right] = 2\mathcal{L}^{-1}\left[\frac{s}{s^2 + 1}\right] + 4\mathcal{L}^{-1}\left[\frac{1}{s^2 + 1}\right] = 2\cos t + 4\sin t,$$

it follows from Theorem A that

$$\mathcal{L}^{-1}\left[\frac{2s}{s^2 - 4s + 5}\right] = \mathcal{L}^{-1}\left[\frac{2s + 4}{s^2 + 1} \Big|_{s \mapsto s-2}\right] = e^{2t} (2\cos t + 4\sin t).$$

In some cases, we employ partial fraction expansion as part of finding the inverse Laplace transform. ■

Example 3:

Find the inverse Laplace transform for each of the functions

$$(a) \frac{8}{s^3 + 4s}$$

$$(b) \frac{3}{s^2 - 4s - 5}$$

$$(c) \frac{8e^{-3s}}{s(s^2 + 4)}$$

The denominator of our function in part (a) is a cubic, whose graph cannot be obtained via a shift of any *quadratic* function. From Calculus, we learn there is a partial fractions expansion of the form

$$\frac{8}{s(s^2 + 4)} = \frac{A}{s} + \frac{Bs + C}{s^2 + 4} = \frac{A(s^2 + 4) + (Bs + C)s}{s(s^2 + 4)} = \frac{(A + B)s^2 + Cs + 4A}{s(s^2 + 4)}$$

Equating coefficients for the various powers of s (and using linear algebra?), we discover that $A = 2$, $B = -2$ and $C = 0$, so

$$\mathcal{L}^{-1}\left[\frac{8}{s^3 + 4s}\right] = \mathcal{L}^{-1}\left[\frac{2}{s} - \frac{2s}{s^2 + 4}\right] = 2\mathcal{L}^{-1}\left[\frac{1}{s}\right] - 2\mathcal{L}^{-1}\left[\frac{s}{s^2 + 4}\right] = 2 - 2\cos(2t).$$

The denominator of the function in part (b) is quadratic, but reducible —i.e., it has real roots, exhibited by the fact that it factors

$$s^2 - 4s - 5 = (s - 5)(s + 1),$$

revealing roots (-1) and 5 . (The quadratic formula would also reveal these *real* roots.) By using partial fraction expansion, we can turn function into the sum of functions with denominators which are 1st degree polynomials:

$$\frac{3}{s^2 - 4s - 5} = \frac{A}{s - 5} + \frac{B}{s + 1} = \frac{A(s + 1) + B(s - 5)}{(s - 1)(s + 5)} = \frac{(A + B)s + (A - 5B)}{(s - 1)(s + 5)}.$$

Equating coefficients of s^1 and s^0 , we can solve to find $A = 1/2$, $B = -1/2$. Thus,

$$\mathcal{L}^{-1}\left[\frac{3}{s^2 - 4s - 5}\right] = \mathcal{L}^{-1}\left[\frac{1/2}{s - 5} - \frac{1/2}{s + 1}\right] = \frac{1}{2}\mathcal{L}^{-1}\left[\frac{1}{s - 5}\right] - \frac{1}{2}\mathcal{L}^{-1}\left[\frac{1}{s - (-1)}\right] = \frac{1}{2}e^{5t} - \frac{1}{2}e^{-t}.$$

The function in part (c) is almost identical to the one in part (a), but for the exponential factor e^{-3s} . (Think Theorem B!) Piggy-backing on our answer to part (a), we obtain

$$\mathcal{L}^{-1}\left[\frac{8e^{-3s}}{s^3 + 4s}\right] = H(t - 3)[2 - 2\cos(2(t - 3))] = 2H(t - 3) - 2H(t - 3)\cos(2(t - 3)).$$

A caution concerning the use of Theorem B to find a Laplace transform


We have noted that Theorems A and B are indispensable when finding inverse Laplace transforms (going from $F(s)$ to $f(t)$), not for the reverse. That is not the same as saying the theorems are not *useful* for finding $F(s)$ from $f(t)$. Look back at Example 1, and check that the theorems provide faster ways of obtaining the answers.

However, it is important to understand that, for a given $f(t)$, Theorem B does *not* address taking the Laplace transform of a "switched on" version of $f(t)$, but rather a "switched on and shifted" version.

Example 4:

Suppose $f(t) = t^2 - 2t + 3$. Then


$$\mathcal{L}[f(t)] = \mathcal{L}[t^2] - 2\mathcal{L}[t] + 3\mathcal{L}[t^0] = \frac{2}{s^3} - \frac{2}{s^2} + \frac{3}{s} = F(s).$$



Theorem B makes it relatively easy to find the Laplace transform of $H(t-1)f(t-1) = H(t-1)[(t-1)^2 - 2(t-1) + 3]$, which has a graph like f but shifted right 1 unit and shifted on at time $t = 1$. By Theorem B,

$$\mathcal{L}[H(t-1)f(t-1)] = e^{-s} \left(\frac{2}{s^3} - \frac{2}{s^2} + \frac{3}{s} \right).$$

The graphs of the time and frequency domain functions appear below.



Since $(t-1)^2 - 2(t-1) + 3 = t^2 - 2t + 1 - 2t + 2 + 3 = t^2 - 4t + 4$, the graph on the left could have been labeled $H(t-1)(t^2 - 4t + 4)$, and the graph on the right is $\mathcal{L}[H(t-1)(t^2 - 4t + 4)]$.

Now, suppose what we desired was actually the Laplace transform of $H(t-1)f(t) = H(t-1)(t^2 - 2t + 3)$, whose graph is depicted at left below. We can only use Theorem B to find

it if we find the formula for the function g for which $g(t-1) = f(t)$; that is,

$$g(t) = f(t+1) = (t+1)^2 - 2(t+1) + 3 = t^2 + 2t + 1 - 2t - 2 + 3 = t^2 + 2,$$


the function obtained shifting f one unit to the *left*. Since

$$\mathcal{L}[g(t)] = \mathcal{L}[t^2 + 2] = \mathcal{L}[t^2] + 2\mathcal{L}[1] = \frac{2}{s^3} + \frac{2}{s},$$

then

$$\mathcal{L}[H(t-1)f(t)] = \mathcal{L}[H(t-1)g(t-1)] = e^{-s} \left(\frac{2}{s^3} + \frac{2}{s} \right),$$

whose graph appears at right below.



Exercises

1. Graph the function and find its Laplace transform.

$$(a) f(t) = t - H(t-1)(t-1) \quad (b) f(t) = H\left(t - \frac{\pi}{4}\right) \cos\left(t - \frac{\pi}{4}\right)$$

$$(c) f(t) = \begin{cases} 0, & t < 3 \\ t^2 + 3t - 8, & t \geq 3 \end{cases} \quad (d) f(t) = \begin{cases} 0, & t < \pi \\ t - \pi, & \pi \leq t < 2\pi \\ 0, & t \geq 2\pi \end{cases}$$

$$(e) f(t) = e^{3t} \sin(4t) \quad (f) f(t) = 4e^{-2(t-5)} H(t-5)(t-5)^2$$

[Note: In the particular case of part (d), you may want to try it both writing it as a series of "shifted, switched-on" functions *and* directly from the definition of Laplace transform, and decide which you think is easier.]

2. Find the inverse Laplace transform for each function.

$$(a) F(s) = \frac{2(s-1)}{s^2 - 2s + 2} \quad (b) F(s) = \frac{2(s-1)e^{-2s}}{s^2 - 2s + 2}$$

$$(c) F(s) = \frac{4}{s^2 - 4} \quad (d) F(s) = \frac{4}{(s-2)^4} + \frac{e^{-2s}}{s^2 + s - 2}$$

$$(e) F(s) = \frac{e^{-s} + e^{-2s} - e^{-3s} - e^{-4s}}{s} \quad (f) F(s) = \frac{s-2}{s^2 - 4s + 3}$$