

Comparing the Growth of Functions as Inputs $(x \text{ or } n) \rightarrow \infty$

Suppose f and g are real-valued functions on a domain that includes nonnegative real numbers. We say that

- f is of order at most g , written $f(x)$ is $O(g(x))$, iff there exists $C > 0$ and $k \geq 0$ such that

$$|f(x)| \leq C|g(x)|, \quad \text{for all real numbers } x > k.$$

We call C, k **witnesses** to this **Big-O** relationship.

- f is of order at least g , written $f(x)$ is $\Omega(g(x))$, iff there exists $C > 0$ and $k \geq 0$ such that

$$|f(x)| \geq C|g(x)|, \quad \text{for all real numbers } x > k.$$

- f is of order g , written $f(x)$ is $\Theta(g(x))$, iff f is simultaneously of order at most g and of order at least g .

Note: Similar definitions hold for sequences (functions from \mathbb{N} to \mathbb{R}).

Examples:

1. Find witnesses that demonstrate $f(x) = 3x^3 + 2x + 7$ is $O(x^3)$.

From graph, it appears $|3x^3 + 2x + 7| \leq 4|x^3|$ when $x > 3$

Or, without graphing:

$$|3x^3 + 2x + 7| = 3x^3 + 2x + 7 \leq 3x^3 + 2x^3 + 7x^3 = 12x^3 = 12|x^3|$$

\uparrow if $x > 0$ \uparrow $x \geq 1$ witnesses $C = 12, k = 1$

got witnesses
 $k = 3, C = 4$

2. Show that $f(x) = \frac{15\sqrt{x}(2x+9)}{x+1}$ is $\Theta(x^{1/2})$.

To show $O(\sqrt{x})$, want a series of \leq starting w/ $\frac{15\sqrt{x}(2x+9)}{x+1}$

$$\frac{15\sqrt{x}(2x+9)}{x+1} \leq \frac{15\sqrt{x}(3x)}{x+1} \leq \frac{15\sqrt{x}(3x)}{x} = 45\sqrt{x}$$

\uparrow if $x \geq 9$


\Rightarrow left fn is $O(\sqrt{x})$ with witnesses $C = 45, k = 9$

Now show f is $\Omega(\sqrt{x})$. Start


$$\frac{15\sqrt{x}(2x+9)}{x+1} \underset{x \geq 0}{\geq} \frac{15\sqrt{x}(2x)}{x+1} \underset{x \geq 1}{\geq} \frac{15\sqrt{x}(2x)}{2x} = 15\sqrt{x}$$


So f is $\Omega(\sqrt{x})$ with witnesses: $C=15, k=1$

Big O: f is $O(g(x))$



f is $\Omega(g(x))$





f is $O(g(x))$ since witnesses C_1, k_1 say so
 f is $\Omega(g(x))$ since witnesses C_2, k_2 say so

$C_1, C_2, \max(k_1, k_2)$ are witnesses
 to f being $\Theta(g(x))$